

Description of η -distributions at RHIC energies in terms of the Ornstein-Uhlenbeck process and relativistic diffusion equations

Minoru BIYAJIMA¹, Masahiro KANEYAMA¹,
Takuya MIZOGUCHI² and Noriaki NAKAJIMA³

¹*Department of Physics, Shinshu University, Matsumoto 390-8621, Japan*

²*Toba National College of Maritime Technology, Toba 517-8501, Japan*

³*Center of Medical Information Science, Kochi University, Kochi, 783-8505 Japan*

Assuming the Ornstein-Uhlenbeck process and relativistic diffusion equations with three sources at $y = 0$, we analyze η -distributions of d +Au collision as well as Au+Au collisions at $\sqrt{s_{NN}} = 130$ GeV and 200 GeV.

1 Introduction: η -distributions in Au+Au collisions at RHIC with various centrality-cuts (C.C.) at RHIC energies have been reported in Refs. 1) and 2). In Ref. 3) we have proposed a stochastic model with the sources at $\pm y_{max}$ and $y = 0$, to describe the η -distributions. That is an extended model from a model of two sources at $\pm y_{max}$ in Refs. 4) and 5) ^{*)}. In this letter, we propose a model with three sources at $y = 0$. A more realistically physical picture is shown in Figs. 1 and 2, where our basic assumption is shown therein.

Very recently PHOBOS Collaboration has reported η -distribution on $dN/d\eta$ ($d + Au$) at $\sqrt{s_{NN}} = 200$ GeV.⁶⁾ Thus we are interested in analyses of the data, to examine whether or not the model proposed in Figs. 1 and 2 works well for descriptions of $dN/d\eta$ (d +Au) as well as $dN/d\eta$ (Au+Au).

2 Theoretical formula: To improve the physical picture of Ref. 3), we consider a formula based on the stochastic approaches. To describe physical picture of Figs. 1 and 2(b), we have found that the following relativistic diffusion equations are available from studies of Ref. 6),

$$\frac{\partial P(y, t)}{\partial t} = \gamma \left[\frac{\partial}{\partial y} (y \pm y_{max}) + \frac{1}{2} \frac{\sigma^2}{\gamma} \frac{\partial^2}{\partial y^2} \right] P(y, t). \quad (1)$$

Using the initial condition

$$P(y, 0) = \delta(y), \quad (2)$$

we obtain the following solutions for the projectile and target fragmentations

$$P^{(T)}(y, y_{max}, t) = \frac{1}{\sqrt{2\pi V_T^2(t)}} \exp \left[-\frac{(y - y_{max}(1 - e^{-\gamma t}))^2}{2V_T^2(t)} \right], \quad (3a)$$

^{*)} In Ref. 5), we have pointed out that single Gaussian distribution with an initial condition $\delta(y)$ cannot explain the whole η -distributions in Au+Au collisions. This fact suggests that contributions from the projectile and target fragmentations are necessary.




Fig. 1. Description of two-step processes by relativistic diffusion equations and the Ornstein-Uhlenbeck process.




Fig. 2. Description of two-step processes for $d+Au$ collisions.

$$P^{(P)}(y, -y_{max}, t) = \frac{1}{\sqrt{2\pi V_P^2(t)}} \exp \left[-\frac{(y + y_{max}(1 - e^{-\gamma t}))^2}{2V_P^2(t)} \right], \quad (3b)$$

where $V_{(P \text{ or } T)}^2 = (\sigma^2/2\gamma)(1 - e^{-2\gamma t})$. For the 3rd source due to gluons in the central region, we assume the Ornstein-Uhlenbeck process (Eq. (1) without the term y_{max} ; see Ref. 3)). The following solution can be derived from Eq. (2) as

$$P^{(C)}(y, t) = \frac{1}{\sqrt{2\pi V_0^2(t)}} \exp \left[-\frac{y^2}{2V_0^2(t)} \right]. \quad (4)$$

Combining Eqs. (3) and (4), we obtain the following expression

$$\begin{aligned} \frac{dn}{d\eta} = J(\eta, m/p_t) \times & \left\{ \frac{c_T}{c_T + c_P + c_0} P^{(T)}(y, y_{max}, t) \right. \\ & + \frac{c_P}{c_T + c_P + c_0} P^{(P)}(y, -y_{max}, t) \Big\} \\ & + J_0 \left\{ \frac{c_0}{c_T + c_P + c_0} P^{(C)}(y, t) \right\}, \end{aligned} \quad (5)$$

where c_T , c_P and c_0 are the weight factors of three contributions for the target fragmentation, the projectile one and the central region, respectively. A relation between y and η , and the Jacobians are given as

$$y = \frac{1}{2} \ln \frac{\sqrt{1 + (m/p_t)^2 + \sinh^2 \eta} + \sinh \eta}{\sqrt{1 + (m/p_t)^2 + \sinh^2 \eta} - \sinh \eta}, \quad (6)$$

$$J(\eta, m/p_t) = \frac{\cosh \eta}{\sqrt{1 + (m/p_t)^2 + \sinh^2 \eta}}, \quad (7)$$

and $J_0 = J(\eta, m_0/p_t)$.

3 η -distributions in Au+Au collisions at smaller centrality-cuts (C.C.): We apply Eq. (5) to the data on η -distributions at the C.C.6-15% and 15-25% at $\sqrt{s_{NN}} = 130$ GeV and 200 GeV by the PHOBOS Collaboration. Our results, using the CERN MINUIT program, are shown in Fig. 3 and Table I. We have to determine the weight factors, c_P , c_T and c_0 . For this aim, we adopt the following criterions $m_0/p_t > \delta(m_0/p_t)$, $m/p_t > \delta(m/p_t)$ and $m_0/p_t < \delta m/p_t$, because pions are mainly produced in the central region. They reveal that the ratio is $c_P : c_0 : c_T = 1 : 6-7 : 1$ at the minimum χ^2 's, where the evolution time $p = (1 - e^{-2\gamma t})$ stops.

Table I. Values of estimated parameters of Eq. (5) with $c_P/c_T = 1$ for Au + Au at $\sqrt{s_{NN}} = 130$ GeV and 200 GeV.

C. C.	130 GeV		200 GeV	
	6-15%	15-25%	6-15%	15-25%
p	0.52±0.21	0.62±0.13	0.78±0.12	0.66±0.16
V_0^2	5.17±0.95	5.98±1.03	5.64±1.42	6.06±1.67
V_T^2	1.31±0.26	1.34±0.30	1.43±0.25	1.59±0.28
c_0	2465±39	1808±32	3078±54	2120±40
c_P	$c_0/6$	$c_0/7$	$c_0/7$	$c_0/6$
m_0/p_t	0.44±0.26	0.37±0.29	0.47±0.33	0.42±0.33
m/p_t	3.60±2.30	2.34±1.06	1.06±0.88	2.06±1.14
$\chi^2/\text{d.o.f.}$	0.64/48	0.82/48	0.50/48	0.54/48

4 η -distributions in d +Au collision: Taking into account the mass ratio, $m_d : m_{Au} = 1 : 100$, we can fix the ratio as $c_T/c_P = 100$. The weight factor c_0 is treated as a free parameter in the present analysis. In eq. (5), two Jacobian factor, $J(\eta, m_T/p_t)$ and $J(\eta, m_P/p_t)$ are used. The evolution parameter $p = 1 - e^{-2\gamma t}$ is also stopped at




Fig. 3. Analyses of data on $dn/d\eta = (N_{ch})^{-1}dN_{ch}/d\eta$ at $\sqrt{s_{NN}} = 130$ GeV and 200 GeV. C.C. = 6-15 % and 15-25 %.

minimum χ^2 . Our result is given in Fig. 4 and Table II. We know that the weight factors are described by $c_P : c_0 : c_T = 0.53 : 40 : 53$. This ratio means contribution from the gluon is the same magnitude of the target fragmentation.

Table II. Parameter values of Eq. (5) with $c_T/c_P = 100$ for $d+Au$ at $\sqrt{s_{NN}} = 200$ GeV.⁶⁾

p	V_0^2	V_P^2	V_T^2	c_0
0.76	4.27 ± 0.64	0.165 ± 0.159	7.56 ± 1.38	39.8 ± 5.4
c_P	m_0/p_t	m_P/p_t	m_T/p_t	$\chi^2/\text{d.o.f.}$
0.529 ± 0.077	0.556 ± 0.166	1.63 ± 0.69	1.88 ± 0.41	$5.1/45$

5 Concluding remarks:

- 1) To improve the previous physical picture in Ref. 3), we assume the relativistic diffusion equations for the target and projectile fragmentations in addition to the Ornstein-Uhlenbeck stochastic process available for that in the central region.
- 2) The ratio, $c_P : c_0 : c_T$, is determined by means of the following criterions;




Fig. 4. Analyses of data on $d+Au$ collision at $\sqrt{s_{NN}} = 200$ GeV by Eq. (5).

$m_0/p_t > \delta(m_0/p_t)$, $m/p_t > \delta(m/p_t)$ and $m_0/p_t < \delta m/p_t$. They can be investigated by analyses of η -distributions with restrictions $\eta_{c1} \lesssim |\eta| \lesssim \eta_{c2}$ in a future.

- 3) As seen in Figs. 3 and 4, our formula seems to be available for the description of η -distributions in $d+Au$ as well as $Au+Au$ collisions.
- 4) From analysis of $d+Au$ collision, $c_P : c_0 : c_T = 0.5 : 40 : 53$, we have know that the contribution of gluon in Au is the same ordered magnitude as that of quarks.
- 5) Since our formula is expressed in the analytical form, it can be applied to other reactions, for example, $Si+Al$ and $S+Au$ collisions. These analyses will be shown elsewhere.⁸⁾

Acknowledgements: We would like to thank R. Nouicer for his kind information on $dn/d\eta$ ($d+Au$), and discussions with N. Suzuki. One of authors (K.M.) is partially supported by Shinshu University.

-
- 1) B. B. Back *et al.* [PHOBOS Collaboration], Phys. Rev. Lett. **87** (2001) 102303.
 - 2) R. Nouicer *et al.* [PHOBOS Collaboration], Proceedings of 37th Rencontres de Moriond on QCD and Hadronic Interactions, Les Arcs, France, 16-23 Mar 2002, arXiv:nucl-ex/0208003.
 - 3) M. Biyajima, M. Ide, M. Kaneyama, T. Mizoguchi and N. Suzuki, Prog. Theor. Phys. Suppl. **153** (2004) 344.
 - 4) M. Biyajima, M. Ide, T. Mizoguchi and N. Suzuki, Prog. Theor. Phys. **108** (2002), 559 [Addendum-ibid. **109** (2003), 151]; See also, M. Ide, M. Biyajima and T. Mizoguchi, arXiv:nucl-th/0302003.
 - 5) M. Biyajima and T. Mizoguchi, Prog. Theor. Phys. **109** (2003) 483.
 - 6) B. B. Back *et al.* [PHOBOS Collaboration], Phys. Rev. Lett. **93** (2004), 082301.
 - 7) G. Wolschin, Phys. Rev. C **69** (2004) 024906; See also, Eur. Phys. J. A **5** (1999), 85.
 - 8) M. Biyajima *et al.*, in preparation.